

AD-A234 601

①

OFFICE OF NAVAL RESEARCH

Contract N00014-84-G-0201

Task No. 0051-865

Technical Report #39

Molecular Orientation of the Silver Tetraneopentoxypthalocyanine and Stearic Acid
Mixed Langmuir-Blodgett Film and Its Chemical and Electrochemical Behaviour

By

Yansong Fu and A.B.P. Lever*

in

Journal of Physical Chemistry

DTIC
ELECTE
APR 19 1991
S B D

York University
Department of Chemistry, 4700 Keele St., North York
Ontario, Canada M3J 1P3

Reproduction in whole, or in part, is permitted for any purpose of the United States Government

*This document has been approved for public release and sale; its distribution is unlimited

*This statement should also appear in item 10 of the Document Control Data-DD form 1473. Copies of the form available from cognizant contract administrator

91 4 17 094

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY Unclassified			3. DISTRIBUTION / AVAILABILITY OF REPORT As it appears on the report		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Report # 39			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION A.B.P. Lever, York University Chemistry Department		6b. OFFICE SYMBOL (if applicable)		7a. NAME OF MONITORING ORGANIZATION Office of Naval Research	
6c. ADDRESS (City, State, and ZIP Code) 4700 Keele St., North York, Ontario M3J 1P3 Canada			7b. ADDRESS (City, State, and ZIP Code) Chemistry Division 800 N. Quincy Street Arlington, VA 22217 U.S.A.		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (if applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-84-G-0201	
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
			WORK UNIT ACCESSION NO		
11. TITLE (Include Security Classification) Molecular Orientation of the Silver Tetraneopentoxypthalocyanine and Stearic Acid Mixed Langmuir-Blodgett Film and Its Chemical and Electrochemical Behavior					
12. PERSONAL AUTHOR(S) Yansong Fu and A.B.P. Lever *					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM Aug. '90 TO Aug. '91		14. DATE OF REPORT (Year, Month, Day) April 4, 1991	
15. PAGE COUNT 30					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Phthalocyanine, Langmuir-Blodgett, Electrochemistry, Polarization, Thin Films		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>Mixed Langmuir-blodgett films comprised of various relative concentrations of silver tetraneopentoxypthalocyanine (AgTNPc) and stearic acid are reported. Electronic spectroscopic polarisation data are discussed in terms of the relative orientation of aggregated AgTNPc domains in the film. Electrochemical data confirm the oxidation of Ag^{II}TNPc(-2) successively to [Ag^{III}TNPc(-2)]⁺ and [Ag^{III}TNPc(-1)]²⁺ in the film phase without loss of orientation. The film was also subject to chemical oxidation and reduction and the degree of organisation following such chemistry is discussed. Ag^{II}TNPc(-2) aggregates, even when diluted with stearic acid, do not display resolved ESR spectra but ternary film mixtures of Ag^{II}TNPc(-2)/H₂TNPc(-2)/stearic acid do exhibit such resolved spectra.</p>					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified/unlimited		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Ronald A. De Marco			22b. TELEPHONE (Include Area Code)		22c. OFFICE SYMBOL

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (2)
Chemistry Division, Code 1113
800 North Quincy Street
Arlington, Virginia 22217-5000

Commanding Officer (1)
Naval Weapons Support Center
Dr. Bernard E. Douda
Crane, Indiana 47522-5050

Dr. Richard W. Drisko (1)
Naval Civil Engineering
Laboratory
Code L52
Port Hueneme, CA 93043

David Taylor Research Center (1)
Dr. Eugene C. Fischer
Annapolis, MD 21402-5067

Dr. James S. Murday (1)
Chemistry Division, Code 6100
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Robert Green, Director (1)
Chemistry Division, Code 385
Naval Weapons Center
China Lake, CA 93555-6001

Chief of Naval Research (1)
Special Assistant for Marine
Corps Matters
Code 00MC
800 North Quincy Street
Arlington, VA 22217-5000

Dr. Bernadette Eichinger (1)
Naval Ship Systems Engineering
Station
Code 053
Philadelphia Naval Base
Philadelphia, PA 19112

Dr. Sachio Yamamoto (1)
Naval Ocean Systems Center
Code 52
San Diego, CA 92152-5000

Dr. Harold H. Singerman (1)
David Taylor Research Center
Code 283
Annapolis, MD 21402-5067

Defense Technical Information Center (2)
Building 5, Cameron Station
Alexandria, VA 22314

ENCLOSURE(2)

147-
ONR Electrochemical Sciences Program
Robert J. Nowak, Program Manager

Professor Hector Abruña
Department of Chemistry
Cornell University
Ithaca, NY 14853
413d018

Professor Allen Bard
Department of Chemistry
The University of Texas at Austin
Austin, TX 78712-1167
413a002

Professor James Brophy
Department of Physics
University of Utah
Salt Lake City, UT 84112
413d015

Professor Bruce Dunn
Department of Materials Science and
Engineering
University of California, Los Angeles
Los Angeles, CA 90024
413d011

Professor Gregory Farrington
Laboratory for Research on the
Structure of Matter
3231 Walnut Street
Philadelphia, PA 19104-6202
413d003

Professor Martha Greenblatt
Department of Chemistry
Rutgers University
Piscataway, NJ 08854
413d008

Professor Adam Heller
Department of Chemical Engineering
University of Texas at Austin
Austin, TX 78712-1062
413h007

Professor C. A. Angell
Arizona State University
Department of Chemistry
Tempe, AZ 85287
413d007

Professor Lesser Blum
Department of Physics
University of Puerto Rico
Rio Piedras, PUERTO RICO 00931
4133002

Professor Daniel Buttry
Department of Chemistry
University of Wyoming
Laramie, WY 82071
4133019

Professor Andrew Ewing
Department of Chemistry
152 Davey Laboratory
Pennsylvania State University
University Park, PA 16802
4133030

Professor W. R. Fawcett
Department of Chemistry
University of California, Davis
Davis, CA 95616
4133020

Professor Joel Harris
Department of Chemistry
University of Utah
Salt Lake City, UT 84112
413a005

Professor Pat Hendra
The University
Southampton SO9 5NH
ENGLAND
4134001

ONR Electrochemical Sciences Program
Robert J. Nowak, Program Manager

Professor Joseph Hupp
Department of Chemistry
Northwestern University
Evanston, IL 60208
4133025

Professor A. B. P. Lever
Department of Chemistry
York University
4700 Keele Street
North York, Ontario M3J 1P3
4131025

Professor Rudolph Marcus
Division of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, CA 91125
4133004

Professor Royce W. Murray
Department of Chemistry
University of North Carolina
Chapel Hill, NC 27514
4133015

Professor Richard Pollard
Department of Chemical Engineering
University of Houston, University Park
4800 Calhoun, Houston, TX 77004
413d016

Dr. Donald Sandstrom
Boeing Aerospace Company
P.O. Box 3999, M/S 87-08
Seattle, WA 98124-2499
4133007

Professor D. E. Irish
Department of Chemistry
University of Waterloo
Waterloo, Ontario, CANADA N2L 3G1
4133017

Professor Nathan S. Lewis
Division of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, CA 91125
413d017

Professor Charles Martin
Department of Chemistry
Texas A&M University
College Station, TX 77843
413d005

Dr. Michael R. Philpott
IBM Research Division
Almaden Research Center
650 Harry Road
San Jose, CA 95120-6099
4133011

Professor B. S. Pons
Department of Chemistry
University of Utah
Salt Lake City, UT 84112
4133010

Professor Jack Simons
Department of Chemistry
University of Utah
Salt Lake City, UT 84112
4131050

ONR Electrochemical Sciences Program
Robert J. Nowak, Program Manager

Dr. H. Gilbert Smith
EG&G Mason Research Institute
57 Union Street
Worcester, MA 01608
413k003

Dr. Stanislaw Szpak
Code 634
Naval Ocean Systems Center
San Diego, CA 92152-5000
4131006

Professor Michael Weaver
Department of Chemistry
Purdue University
West Lafayette, IN 49707
4133001

Professor Geroge Wilson
Department of Chemistry
University of Kansas
Lawrence, KS 66045
413k002

Professor Ernest Yeager
Case Center for Electrochemical
Sciences
Case Western Reserve University
Cleveland, OH 44106
4133008

Professor Ulrich Stimming
Department of Chemical Engineering
and Applied Chemistry
Columbia University
New York, NY 10027
4133014

Professor Petr Vanýsek
Department of Chemistry
Northern Illinois University
DeKalb, IL 60115
413k001

Professor Henry White
Department of Chemical Engineering
and Materials Science
421 Washington Ave., SE
Minneapolis, MN 55455
400o027yip

Professor Mark S. Wrighton
Department of Chemistry
Massachusetts Institute of Technology
Cambridge, MA 02139
4131027

Dept. of Chemistry, York University,
North York, Toronto, Ontario, Canada, M3J 1P3

**Molecular Orientation of the Silver Tetraneopentoxypthalocyanine and Stearic
Acid Mixed Langmuir-Blodgett Film and Its Chemical and Electrochemical Behavior**

by Yansong Fu and A.B.P. Lever*

Abstract

Mixed Langmuir-blodgett films comprised of various relative concentrations of silver tetraneopentoxypthalocyanine (AgTNPc) and stearic acid are reported. Electronic spectroscopic polarisation data are discussed in terms of the relative orientation of aggregated AgTNPc domains in the film. Electrochemical data confirm the oxidation of $\text{Ag}^{\text{II}}\text{TNPc}(-2)$ successively to $[\text{Ag}^{\text{III}}\text{TNPc}(-2)]^+$ and $[\text{Ag}^{\text{III}}\text{TNPc}(-1)]^{2+}$ in the film phase without loss of orientation. The film was also subject to chemical oxidation and reduction and the degree of organisation following such chemistry is discussed. $\text{Ag}^{\text{II}}\text{TNPc}(-2)$ aggregates, even when diluted with stearic acid, do not display resolved ESR spectra but ternary film mixtures of $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{H}_2\text{TNPc}(-2)/\text{stearic acid}$ do exhibit such resolved spectra.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Introduction

The phthalocyanines,^{1,2} known for some 60 years, and used primarily in the dye, paint and ink industry, now seem ready for exploitation in a wide range of other diverse applications.³ The Langmuir-Blodgett (LB) technique for generating monolayer structures has considerable promise for commercial applications involving these phthalocyanine species.⁴ Such applications include electrochromic devices,⁵ molecular electronics^{6,7}, switching devices⁸ and, especially, chemical gas sensors.⁹⁻¹² Success is largely dependent on the degree of the molecular organization within the film and the ability to control and reproduce such organization. Recently, Vandevyver and Barraud¹³ have reviewed the techniques for the LB film characterization, among which linear dichroism measurement,^{14,15} grazing incidence reflection infrared spectroscopy¹⁶ and ESR spectroscopy^{17,18} are the most convenient and commonly available methods to investigate molecular orientation in the film.

We have recently described¹⁹ the preparation and characterization of a new soluble silver^{II} phthalocyanine, Ag^{II}TNPc (TNPc = tetraneopentoxypthalocyanine). The solubility in a range of non-donor organic solvents provides a means of studying silver phthalocyanine in a fashion unavailable previously for the insoluble unsubstituted analogue. The neopentoxy groups are randomly distributed among the 3 and 4 positions of the outer benzene rings so that this species is isolated as a mixture of geometric isomers which are not separable by the usual chromatographic procedures. Nevertheless it has proven possible to generate organized films therewith.

The present study describes the formation of a mixed LB film of Ag^{II}TNPc and stearic acid which is characterized using dichroism measurements to show the average molecular orientation of the film. Strong intermolecular aggregation is present as monitored by UV-Vis and ESR spectra.

While it is sometimes difficult to transfer a pure phthalocyanine monolayer onto a solid substrate, a mixture of the phthalocyanine with some long chain amphiphilic

species, such as octadecanol,²⁰ stearic²¹⁻²³ or arichidic acid,²⁴ has proven effective. It would be interesting to know how the long chain species such as stearic acid functions in the film, and how this might affect the molecular orientation and the chemistry of the co-partner. The understanding of this issue is important to the utility of those long chain hydrocarbon species in making mixed LB films.

To further characterize the film, linear cyclic voltammetry (LCV) and differential pulse cyclic voltammetry (DPV) were carried out on the film deposited on a SnO_2 film-coated glass slide. The stability of the molecular organization within the film was also investigated by monitoring the change in dichroic ratio of the film upon electrochemical or chemical modification.

Experimental Details

Materials: Metal-free tetraneopentoxyl phthalocyanine (H_2TNPc) was prepared following the literature methods.²⁵ Silver^{II} tetraneopentoxypthalocyanine ($\text{Ag}^{\text{II}}\text{TNPc}$) was prepared according to the recently described method.¹⁹ Toluene, as spreading solvent, was Aldrich HPLC grade and glass distilled. The subphase was water purified by double distillation over KMnO_4 followed by passage through a Barnstead organic removal cartridge and two Barnstead mixed resin ultrapure cartridges.

Spectroscopic and Electrochemical Methods: The film electronic spectra and dichroic ratio were measured with a Hitachi Perkin - Elmer Microprocessor 340 spectrometer. Polaroid film polarizers were mostly employed, but some late experiments used calcite polarizers and a Cary 2415 spectrometer. A Varian E4 EPR spectrometer was used to record ESR spectra. Cyclic and differential pulse voltammetry were performed with a Princeton Applied Research (PARC) model 173 Potentiostat/Galvanostat, controlled by a PARC 175 Universal Programmer, and a PARC 174A Polarographic Analyzer. Spectro-electrochemical data were collected with a Guided Wave Inc. model 100 Spectrum Analyzer and the PARC model 173 Potentiostat.

Preparation of the LB Film: The LB trough was built at York following a described design.²⁶ The measurements of surface pressure (π) versus area per molecule (A) isotherms and multilayer deposition were carried out with an Apple II+ microcomputer controlled LB system, using a local program. In general, a monolayer was compressed at a speed of $1.5 \text{ \AA}^2/\text{molecule}/\text{min.}$ and then transferred onto a substrate at $5 \text{ mm}/\text{min.}$ Before the actual deposition, about 10 minutes are allowed for the self-organization of the molecules after forming the monolayer. Surface pressure was measured based on Wilhelmy's principle,²⁷ calibrated with a stearic acid monolayer, whose collapse pressure was reported to be $40 \text{ mN}/\text{M.}$ ²⁷ The spreading solution was usually of $8 \times 10^{-4} \text{ M}$ and the deposition pressure was about $20 \text{ mN}/\text{M.}$

Glass and quartz slides were first treated with boiling CH_2Cl_2 or CHCl_3 and then sonicated in a 1 M NaOH aqueous solution. The resulting hydrophilic surface of the slide was made hydrophobic by immersing the slide in a $5\% \text{ (Vol.) Me}_2\text{Cl}_2\text{Si}/\text{CCl}_4$ (Me - methyl) solution for a few minutes. For SnO_2 film-coated glass slides, a slightly basic Decon solution was used instead of the NaOH solution. A carefully stored, cleaned slide is usable for a few days before re-cleaning would be necessary.

Results and Discussions

Pure Ag^{II} TNPc Monolayer: Figure 1 (curve a) shows the surface pressure (π) versus area per molecule isotherm for a pure Ag^{II} TNPc monolayer. The monolayer has a fairly large compressibility, as implied by the fact that the surface pressure begins to rise at about 90 \AA^2 and collapses at about 55 \AA^2 , giving an area of 35 \AA^2 for the compression. The large compressibility is attributed to the interaction between the neopentoxyl groups of the molecule. It can also be seen that the monolayer collapses at a relatively low pressure, around $16 \text{ mN}/\text{M}$, suggesting that a fragile monolayer has been formed on the subphase. The fragility of the film led to a poor deposition ratio (the area loss on the subphase surface over the area coated on the solid substrate), being

only 0.2 or so, and thus, no useful film of pure $\text{Ag}^{\text{II}}\text{TNPc}$ was obtained. The low collapse pressure of the $\text{Ag}^{\text{II}}\text{TNPc}$ monolayer is due to the large compressibility causing the monolayer to fold easily. In addition, the area per molecule of the monolayer at zero surface pressure was found to be 76 \AA^2 , indicating a tilted orientation of the phthalocyanine macrocyclic plane with respect to the subphase surface.

Mixed Films of $\text{Ag}^{\text{II}}\text{TNPc}$ and Stearic Acid: When a single-component monolayer shows a poor deposition property, improved behaviour can often be observed by mixing the pure monolayer with some long chain hydrocarbon alcohol or acid. In our study, stearic acid ($\text{C}_{18}\text{H}_{36}\text{O}_2$, SteA) was mixed with the $\text{Ag}^{\text{II}}\text{TNPc}$ species to improve the deposition ratio. The pressure isotherm of the $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ mixed monolayer with a molar ratio of $1:2 = \text{Ag}^{\text{II}}\text{TNPc}:\text{SteA}$ is also shown in Figure 1 (curve d). Compared to curve (a) in the same figure, the mixed monolayer showed a much smaller compressibility and a higher collapse pressure (26 mN/M), indicating that a more compact monolayer had been obtained.

The mixed monolayer can be readily deposited onto hydrophobic surfaces of different kinds of solid substrates, such as a glass/quartz slide, SnO_2 film-coated glass slide, and an HOPG (highly-oriented pyrolytic graphite) plate, all with a deposition ratio of nearly unity. The total molecular area occupied by the mixed monolayer on the water surface equals the sum of the individual molecular areas of the two components within experimental error when assuming the molecular areas to be 76 \AA^2 for $\text{Ag}^{\text{II}}\text{TNPc}$ and 21 \AA^2 for SteA. This observation suggests that the $\text{Ag}^{\text{II}}\text{TNPc}$ molecules and SteA molecules all have direct contact with subphase surface, indicative of immiscibility of the $\text{Ag}^{\text{II}}\text{TNPc}$ and stearic acid components.

The $\text{Ag}^{\text{II}}\text{TNPc}$ molecules are believed to be packed tightly by the SteA molecules; when a hydrophobic substrate is dipped into the water, its surface interacts with the

SteA hydrocarbon long chain strongly enough to bring the entire mixed monolayer onto the substrate. X-type transference (the monolayer can only be transferred upon the downward stroke of a substrate) was observed, which probably suggests that after the first layer was transferred, the deposition of the successive layers involved only the interaction between the neopentoxyl groups of the $\text{Ag}^{\text{II}}\text{TNPc}$ and the hydrocarbon long chains of the SteA.

In Figure 2, the electronic spectrum (curve a) of the mixed film displays a normal Soret band at 350 nm and a Q-band at about 620 nm with a shoulder at about 680 nm. The 620 nm Q-band is blue shifted from the corresponding band in solution,¹⁹ indicative of strong inter-molecular aggregation;^{23,28,29} the 680 nm shoulder probably indicates a small amount of monomer species exists in the film, though it could arise from transitions within low symmetry aggregated species.²⁹

One of the $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ mixed films was washed off the glass slide with toluene and the spectrum of the resultant solution (about 10^{-5}M) was taken, as shown in Figure 2-b. The spectrum looks identical to that of a newly-prepared $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{toluene}$ solution,¹⁹ suggesting that no chemical change has occurred to the $\text{Ag}^{\text{II}}\text{TNPc}$ molecules.

The inset in Figure 3, an ESR spectrum taken from an $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ (molar ratio, 1:2) mixed film, shows only a single broad peak arising from the unpaired electron in the Ag^{II} ion. The strong inter-molecular interaction has obscured the hyperfine coupling due to the silver nucleus and the four nitrogen nuclei, which otherwise should yield eleven peaks.^{19,30}

However, using a ternary film composed of $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{H}_2\text{TNPc}(-2)/\text{SteA}$ (1:11:24), hyperfine coupling was observed with anisotropic g values (Figure 3). Thus in this case, the H_2TNPc molecules are able to slip between adjacent $\text{Ag}^{\text{II}}\text{TNPc}$ molecules and reduce the degree of inter- Ag^{II} relaxation.

Dichroic Ratio Measurements: When polarized light is used to record the electronic spectrum of a well-organized film, the absorption of the light is expected to vary with the angle between the direction of the electric vector of the light and the dipping direction when making the film. Shown in Figure 4 are the electronic absorption spectra of two $\text{Ag}^{\text{II}}\text{TNPC}(-2)/\text{SteA}$ mixed films with the electric vector perpendicular (E_{\perp}) and parallel (E_{\parallel}) to the dipping direction. The fact that Abs_{\perp} is larger than Abs_{\parallel} suggests that most of the $\text{Ag}^{\text{II}}\text{TNPC}$ molecules in the film tend to sit on the substrate surface with their macrocyclic plane facing the dipping direction. Moreover since this film is 12 layers thick, the films must lay down on top of one another in a well organised fashion.

A series of $\text{Ag}^{\text{II}}\text{TNPC}(-2)/\text{SteA}$ mixtures have been investigated with the molar ratios being 2:1, 1:1, 1:2, 1:4, and 1:9, in order to determine how the SteA molecules affect the deposition of the $\text{Ag}^{\text{II}}\text{TNPC}$ monolayer and their molecular orientation. The π -A isotherms of the five mixtures are shown in Figure 1. These illustrate that the collapse pressure of the mixed monolayer increases with the amount of SteA in the mixture; the more SteA a monolayer contains, the more difficult it will be to fold the monolayer. The experiment has also shown that when the $\text{Ag}^{\text{II}}\text{TNPC}(-2)/\text{SteA}$ molar ratio was larger than one, the deposition ratio became less than unity, with the 2:1 mixture only exhibiting a value of 0.5. This could be understood by recalling that the pure $\text{Ag}^{\text{II}}\text{TNPC}$ monolayer can only be transferred with difficulty.

Each film shows a dichroic effect but to a varying degree. The film with a higher $\text{Ag}^{\text{II}}\text{TNPC}(-2)/\text{SteA}$ molar ratio gave a higher value of the dichroic ratio ($\text{Abs}_{\perp}/\text{Abs}_{\parallel}$), indicative of a higher degree of molecular organization. Two examples are shown in Figure 4 with (a) from the 2:1 mixed film and (b) from the 1:4 mixed film, and in Figure 5 the dichroic ratios for five mixtures are plotted versus the molar fraction of $\text{Ag}^{\text{II}}\text{TNPC}$ in the mixture. This clearly illustrates that the molecules in the mixed film become more organized with an increase in the relative concentration of

$\text{Ag}^{\text{II}}\text{TNPc}$. Thus the SteA molecules do not help to organize the $\text{Ag}^{\text{II}}\text{TNPc}$ molecules but instead reduce the orderliness of the film.

Previous studies with stearic acid diluted phthalocyanine films,²³ have demonstrated the existence of aggregated domains within the stearic acid arrays. Our data suggest a similar situation. Individual aggregates show high organisation and dichroism, and at low stearic acid concentrations, neighboring domains may serve to orient each other in a parallel fashion. As the relative concentration of phthalocyanine is reduced, the domains are further apart and their orientation more random; thus the dichroism is reduced.

To gain further insight into the molecular orientation in the film, two angles, θ and ϕ , are proposed to represent the average molecular orientation. This is shown schematically in Figure 6, in which Y corresponds to the dipping direction when making the film, Z is the normal of the substrate surface and X is perpendicular to the YZ plane. If a flat circle is used to represent the phthalocyanine transition dipole plane, the angle formed by the normal (z') to the circle with respect to the Z axis will be called θ , and the angle between the projection of the normal in the XY plane and the X axis called ϕ .

Following the procedure developed by Yoneyama et al.,³¹ orientation angles, θ and ϕ , for each of the five mixed films can be calculated and the results are listed in Table 1. There is no consistent change in the θ value, but the value of ϕ decreases with increasing relative concentration of SteA. Evidently the rotation of $\text{Ag}^{\text{II}}\text{TNPc}$ molecules about their x' axis, which is parallel to the X axis, is not affected by the presence of SteA, but the rotation about the r' axis, which is perpendicular to the normal of the phthalocyanine circle and parallel to the YZ plane, will be easier if the amount of SteA is increased. Considering that a film with randomly ordered molecules should have an average ϕ value of 45° ($\int_0^{90^\circ} \phi d\phi / \int_0^{90^\circ} d\phi = 45^\circ$) and a perfectly ordered film should show 90° for ϕ , Table 1 also suggests that the molecular organization in the

film becomes worse in SteA mixed films.

In summary, the above observations reveal that although the presence of SteA is necessary to obtain uniform films, the molecular organization of the film is reduced by giving more freedom of orientation for the $\text{Ag}^{\text{II}}\text{TNPc}$ molecules.

Cyclic Voltammetric Studies: An $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ (1:2) mixed film (3 layers thick) was deposited onto a SnO_2 film-coated glass slide, and studied by cyclic voltammetry in a 0.2M NaClO_4 aqueous solution. The film voltammogram exhibited a broad and irreversible redox wave with the half-potential, $E_{1/2} = 0.65\text{V}$ (vs. SCE), as shown in Figure 7a. The redox process is a one-electron oxidation, leading to the species $[\text{Ag}^{\text{III}}\text{TNPc}(-2)]^+$ based upon the following arguments.

(1) In the solution voltammetry, $\text{Ag}^{\text{II}}\text{TNPc}$ is successively oxidized in two one-electron steps to $[\text{Ag}^{\text{III}}\text{TNPc}(-2)]^+$ and $[\text{Ag}^{\text{III}}\text{TNPc}(-1)]^{2+}$ at 0.71 and 1.19 V (vs SCE) respectively.¹⁹

(2) When the same kind of film, deposited on a glass slide, was oxidized chemically with chlorine gas, oxidation occurred to yield $[\text{Ag}^{\text{III}}\text{TNPc}(-2)]^+$, as indicated by electronic spectroscopy (vide infra).

(3) A spectroelectrochemical study of the film polarized at +0.75V showed a spectrum with a red shifted and slightly less intense Q-band and a new band around 440 nm (Figure 8). This is consistent with the solution spectroelectrochemical data wherein the formation of $[\text{Ag}^{\text{III}}\text{TNPc}(-2)]^+$, from $\text{Ag}^{\text{II}}\text{TNPc}$ in DCB, is associated with a red shifted and less intense Q band, and the appearance of a new absorption at 430 nm.¹⁹ Upon re-reduction of the oxidized film by polarization at -0.05 V, the 440 nm band disappeared, and the Q band shifted back to its original position with increased intensity. This cycle can be repeated many times, but with some loss of overall Q band intensity each time, probably due to some film loss from the electrode. These observations are inconsistent with the postulate that polarization merely changes the aggregate-monomer

ratio, since such a change towards the monomer would result in an increase in intensity of the red shifted Q band.

In the solution electrochemistry, the first oxidation step of $\text{Ag}^{\text{II}}\text{TNPc}(-2)$ to $[\text{Ag}^{\text{III}}\text{TNPc}(-2)]^+$ is very broad and ill-resolved due to extensive aggregation and a kinetically slow electron transfer step. The $\text{Ag}^{\text{II}}\text{TNPc}(-2)$ LB film is also greatly aggregated, as discussed above. This molecular interaction also results in a broad CV peak for the LB film whose irreversibility is indicated by the large peak separation ($E_a - E_c$), of over 200mv. The differential pulse cyclic voltammogram, (in Figure 7b), shows that the peak is much smaller upon reversing the scanning. This may be associated with the movement of ClO_4^- anions through the film. Upon oxidation of the film, the ClO_4^- anions penetrate into the film to balance the positive charge. However, after their penetration, the anions are locked inside the film and have difficulty leaving on the time scale of the experiment. The scan rate dependence of the redox wave shows a nonlinear relationship between peak current and scan rate, consistent with this supposition.

Chemical Stability of the LB Film: An important property for an LB film is the stability of the molecular organization of the film against electrochemical or chemical change. We have therefore studied the change in dichroic ratio of the LB film upon electrochemical or chemical modification.

The polarized UV-Vis spectra of a $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ (1:2) mixed LB film (6 layers) deposited on a SnO_2 -film coated glass slide is shown in Figure 9a. They resemble the spectra of the film on a regular glass slide, except for a small broad band around 500 nm, which is believed to be due to a specific interaction with the SnO_2 surface since the absorption remains if the film is oxidized or reduced electrochemically, but is not present in the solution spectrum of the film washed off the surface.

Electrochemical studies were carried out under the same conditions as those employed in the cyclic voltammetric studies. After repeating the cyclic potential scanning in the range of -0.4 to 1.0 V for 9 times, the polarized spectra were recorded (Figure 9b). The dichroic ratio when measured at the main Q-band measured close to 100 % of its original value, although the interaction absorption at 500 nm had increased. The invariance of the dichroic ratio suggests that the molecular organization within the film is stable in terms of general cyclic voltammetric measurement.

Chemical Cycling of LB Film: An $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ (1:2) film (10 layers) deposited on a glass slide was treated with molecular Cl_2 gas either in the gas or aqueous phase. In the gas phase experiment, the slide was placed, for a few minutes, inside a bottle containing Cl_2 gas, approximately 0.5% by volume in air, and then removed and washed with distilled water. The resulting spectrum is shown in Figure 10b. Compared to the original film spectrum (Figure 10a), the Cl_2 -treated film showed a red-shifted Q-band at 640 nm but no new band around 500 nm, consistent with the one-electron oxidation of $\text{Ag}^{\text{II}}\text{TNPc}(-2)$ to $[\text{Ag}^{\text{III}}\text{TNPc}(-2)]^+$. The Cl_2 -treated film maintained the same degree of molecular organization as before, indicated by the almost unchanged dichroic ratio (1.23 in Figure 10a, and 1.19 in Figure 10b).

When the gas phase Cl_2 -oxidized film was immersed in a 10% (Vol) N_2H_4 aqueous solution for 15 to 30 min., the initial Q band envelope was restored entirely with a slight decrease in dichroic ratio (to 1.17), as shown in Figure 10c. Thus the LB film can be oxidized by chlorine gas and reduced back with aqueous hydrazine without change in film organization. The spectra of the $[\text{Ag}^{\text{III}}\text{TNPc}(-2)]^+$ species in Figures 8 and 10 likely differ because the counter-ions and, possibly, coordination number are different in the two cases.

In a solution phase oxidation, the same film as above was exposed to a 0.1 M NaClO_4 aqueous solution bubbled with chlorine gas (10 - 20 s) for 2 minutes. In this

case, the electronic spectrum again demonstrated oxidation to $[\text{Ag}^{\text{III}}\text{TNPc}(-2)]^+$ but with a slightly larger red shift in the Q band perhaps because the reaction was more complete. However the dichroic ratio was greatly reduced (Figure 11a), implying substantial loss of film organization. When this film was reduced back to $\text{Ag}^{\text{II}}\text{TNPc}$ (Figure 11b) with hydrazine, the dichroic ratio declined almost to unity, indicating loss of order. A much thinner film (4 layers) was used to repeat the liquid phase experiment, with the same results being obtained.

Thin film reorganization upon redox sampling undoubtedly involves solvent transfer through the film.^{32,33} Thus the loss of organization upon solvent phase oxidation, but retention during gas phase oxidation may reflect the disruptive effect of such solvent transport in the former case.

Summary and Conclusions: Highly ordered well behaved $\text{Ag}^{\text{II}}\text{TNPc}(-2)$ Langmuir-Blodgett films can be co-deposited with stearic acid onto suitable substrates, with a deposition ratio of unity. Increasing stearic acid facilitates the transfer of a compact film, but decreases the degree of organization of the $\text{Ag}^{\text{II}}\text{TNPc}(-2)$ component. The ESR spectra of the mixed $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ films show that aggregation is still occurring, evidence that the stearic acid does not interleave between the $\text{Ag}^{\text{II}}\text{TNPc}(-2)$ molecules. Thus dilute films are likely composed of domains of aggregated $\text{Ag}^{\text{II}}\text{TNPc}(-2)$ which are internally ordered, separated by stearic acid, but with some loss of relative organization between the domains. The presence of the dichroism in the film spectra suggests that each domain consists of $\text{Ag}^{\text{II}}\text{TNPc}(-2)$ molecules with their macrocyclic plane tending to face the dipping direction. In the films of higher relative $\text{Ag}^{\text{II}}\text{TNPc}$ concentration, each of the domains has a similar orientation, resulting in the anisotropic optical property. The degree of freedom with which the domains orient is dependent upon the relative amount of SteA in the mixture. A larger relative amount of SteA results in a larger degree of freedom for domain

orientation and, therefore, worsens the film organization.

Film organization is largely unaffected by electrochemical cycling or by chemical cycling using gaseous chlorine and aqueous hydrazine, but the order is rapidly lost if the film is cycled using a solution chemical oxidant, perhaps due to solvent transport. Retention during electrochemical oxidation in solution shows that different solvent transport and likely anion compensation pathways occur depending upon whether the method of oxidation is chemical or electrochemical. Clearly the detailed mechanisms require further study.

Acknowledgements We are indebted to the Natural Sciences and Engineering Research Council (Ottawa), and the Office of Naval Research (Washington) for financial support.

Reference

- (1) Lever, A. B. P. Adv. Inorganic Chem. and Radiochem 1965, 7, 27.
- (2) Leznoff, C. C.; Lever, A. B. P. Eds. Phthalocyanines. Properties and Applications; VCH: New York, 1989.
- (3) Lever, A. B. P. CHEMTECH 1987, 17, 506.
- (4) Fourth International Conference on Langmuir-Blodgett Films, Tsukuba, Japan, April 1989, Thin solid Films 1989, 179, 1-554.
- (5) Yamamoto, H.; Sugiyama, T.; Tanaka, M. Japn. J. Appl. Phys. 1985, 24, L 305.
- (6) Sugi, M. J. Molecular Electronics 1985, 1, 3.
- (7) Hann, R. A.; Gupta, S. K.; Fryer, J. R.; Eyres, B. L. Thin Solid Films 1985, 134, 35.
- (8) Vandevyver, M.; Ruaudel-Teixier, A.; Barraud, A. J. Chim. Phys. 1985, 82(6), 707.
- (9) Tredgold, R. H.; Young, M. C. J.; Hodge, P.; Hoorfar, A. IEEE Proc. J. 1985, 132, 152.
- (10) Fujiki, M.; Tabei, H.; Imamura, S. Japn. J. Appl. Phys. 1987, 26, 1224.
- (11) Reswick, R. B.; Pitt, C. W. Chem. Phys. Letters 1988, 143, 589.
- (12) Palacin, S.; Ruaudel-Teixier, A.; Barraud, A. J. Phys. Chem. 1986, 90, 6237.
- (13) Vandervyver M.; Barraud, A. J. Molecular Electronics 1988, 4, 207.
- (14) Kalina, D. W.; Crane, S. W. Thin Solid Films 1985, 134, 109.
- (15) Ogawa, K.; Kinoshita, S.; Yonehara, H.; Nakahara, H.; Fukuda, K. J. Chem. Soc., Chem. Commun 1989, 477.

- (16) Mumby, S. J.; Swalen, J. D.; Rabolt, J. F. Macromolecules **1986**, 19, 1054.
- (17) Cook, M. J.; Daniel, M. F.; Dunn, A. J.; Gold, A. A.; Thomson, A. J. J. Chem. Soc., Chem. Commun. **1986**, 863.
- (18) Pace, M. D.; Barger, W. R.; Snow, A. W. J. Mag. Res. **1987**, 75, 73.
- (19) a) Fu, G.; Fu, Y.; Lever, A. B. P. Inorg. Chem. **1990**, 29, 4090. b) Ouyang, J.; Lever, A. B. P. J. Phys. Chem. **1991**, 95, in press. c) Ouyang, J.; Lever, A. B. P. J. Phys. Chem. **1991**, 95, accepted.
- (20) Barger, W. R.; Snow, A. W.; Wohltjen, H.; Jarvis, N. L. Thin Solid Films **1985**, 133, 197.
- (21) Palacin, S.; Ruaudel-Teixier, A.; Barraud, A. J. Phys. Chem. **1989**, 93, 7195.
- (22) Watanabe, I.; Hong, K.; Rubner, M. F. J. Chem. Soc., Chem. Commun **1989**, 123.
- (23) Petty, M.; Lovett, D. R.; O'Connor, J. M.; Silver, J. Thin Solid Films **1989**, 179, 387.
- (24) Aroca, R.; Battisti, D. Langmuir **1990**, 6, 250.
- (25) Leznoff, C. C.; Greenberg, S.; Marcuccio, S. M.; Minor, P. C.; Seymour, P.; Lever, A. B. P.; Tomer, K. B. Inorg. Chim. Acta **1984**, 89, L35.
- (26) Van Ryswyk, H.; Eisenhart, J. M.; Blom, N. Rev. Sci. Instrum. **1985**, 56, 2339.
- (27) Gaines, Jr., G. L. Insoluble Monolayers At Liquid-Gas Interface; Interscience: New York, **1966**.
- (28) Snow, A. W.; Jarvis, N. L. J. Am. Chem. Soc. **1984**, 106, 4706.
- (29) Dodsworth, E. S.; Lever, A. B. P.; Seymour, P.; Leznoff, C. C. J. Phys. Chem. **1985**, 89, 5698.
- (30) Kneubuhl, F. K.; Koski, W. S.; Caughey, W. S. J. Am. Chem. Soc. **1961**,

83, 1607.

(31) Yoneyama, M.; Sugi, M.; Saito, M. Japn. J. Appl. Phys. **1986**, 25, 961.

(32) Bruckenstein, S.; Hillman, A. R.; J. Chem. Phys. **1988**, 92, 4837.

(33) Bruckenstein, S.; Wilde, C. P.; Hillman, A. R.; J. Chem. Phys. **1990**,
94, 6458.

Table 1 Molecular Orientation as a Function of Molar Fraction

Mol.Frac.		
Ag ^{II} TNPc	θ	ϕ

0.10	52	46
0.20	52	53
0.33	42	63
0.50	46	72
0.67	48	79

See text and Figure 6 for coordinate system.

Figure Legends

Figure 1. Surface pressure vs. area per molecule isotherms for a) pure $\text{Ag}^{\text{II}}\text{TNPc}$, and the five $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ mixtures with molar ratios being ($\text{Ag}^{\text{II}}\text{TNPc}:\text{SteA}$): b) 2:1, c) 1:1, d) 1:2, e) 1:4, f) 1:9.

Figure 2. Electronic absorption spectra of a) $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ (1:2) mixed LB film, with a thickness of 12 layers, and b) $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{toluene}$ solution, about $5 \times 10^{-5} \text{ M}$, path length 0.2 cm, made by washing off the LB film with toluene.

Figure 3. ESR spectra of $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{H}_2\text{TNPc}(-2)/\text{SteA}$ (1:11:24) mixed LB film of 40 layers on quartz slide. The inset in the figure: $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ (1:2) mixed LB film of 50 layers on quartz slide.

Figure 4. Polarized electronic spectra of $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ mixed LB film (12 layers thick) with molar ratios ($\text{Ag}^{\text{II}}\text{TNPc}:\text{SteA}$) being a) 2:1 and b) 1:4, and the electric vector of the incident light being α) perpendicular (E_{\perp}) and β) parallel (E_{\parallel}) to the dipping direction.

Figure 5. A relationship between dichroic ratios of $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ mixed LB films and the molar fraction of $\text{Ag}^{\text{II}}\text{TNPc}$.

Figure 6. Coordinate system for showing the molecular orientation of $\text{Ag}^{\text{II}}\text{TNPc}$ in the film. Y is the dipping direction, Z is the normal of the substrate surface and X is perpendicular to the YZ plane.


Figure 7. a) Linear cyclic voltammogram (scan rate 100 mV/s) and b) differential pulse cyclic voltammogram (scan rate 2 mV/s) of $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ (1:2) mixed LB film on a SnO_2 film-coated glass slide (8 layers thick) in 0.2M NaClO_4 aqueous solution.


Figure 8. Electronic spectra of $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ (1:1) mixed LB film (12 layers thick) and the oxidized species generated by potentiostatic electrolysis, the potential was set at 0.75V vs. SCE.

Figure 9. Dichroic measurements of $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ (1:2) mixed LB film (6 layers) on a SnO_2 film-coated glass slide with a) before and b) after cyclic voltammetric studies (9 cycles).


Figure 10. Polarized spectra of $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ (1:2) mixed LB film (10 layers) on glass slide upon chemical reactions: a) original film; b) 5 minute exposure to 0.5%(Vol.) Cl_2 gas in air; c) 25 minute immersion of the Cl_2 -treated film in a 10%(Vol.) N_2H_4 aqueous solution.



Figure 11. Polarized spectra of $\text{Ag}^{\text{II}}\text{TNPc}(-2)/\text{SteA}$ (1:2) mixed LB film (10 layers) on glass slide. a) The film has been treated with a Cl_2 -bubbled (15 secs) 0.1M NaClO_4 aqueous solution for 3 minutes; b) the oxidized film was treated with a 10%(Vol.) N_2H_4 aqueous solution for 25 minutes.






222 Fig










Fig. 1

